

## Anti-K<sub>Ca</sub>2.3 (C-term) (SK3)

(Small conductance Ca<sup>2+</sup>-activated K<sup>+</sup> Channel 3, SK3, SKCa3, KCNN3)

**Product #:** APC-103  
**Current Lot #:** AN-01  
**Host:** Rabbit  
**Type:** Polyclonal.

**Epitope:** Peptide (C)KQIGSLESKLEHLTAS, corresponding to amino acid residues 659-674 of human K<sub>Ca</sub>2.3 (Accession [Q9UGI6](#)).

**Epitope location:** Intracellular, C-terminal part.

**Homology with other species:** Rat, mouse, pig-identical.

**Reactivity Confirmed:** Rat, human.

**Identity of the peptide:** Confirmed by mass-spectrography and amino acid analysis.

**Purity:** Affinity purified on immobilized antigen.

**Sample Sizes:** 50 µl or 0.2 ml after reconstitution of the lyophilized powder.

**Storage before Reconstitution:** Lyophilized powder can be stored intact at room temperature for several weeks. For longer periods, it should be stored at -20°C.

**Reconstitution:** 50 µl or 0.2 ml deionized water, depending on the sample size.

**Antibody Concentration after Reconstitution:** 0.8 mg/ml.

**Buffer after Reconstitution:** Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.025% NaN<sub>3</sub>.

**Storage after Reconstitution:** The reconstituted solution can be stored at 4°C for up to 2 weeks. For longer periods, small aliquots should be stored at -20°C or below. Avoid multiple freezing and thawing. Further dilutions should be made using a carrier protein such as BSA (1%). Centrifuge **all** antibody preparations before use (10000 × g 5 min).

**Control Antigen Storage before Reconstitution:** Lyophilized powder can be stored intact at room temperature for several weeks. For longer periods, it should be stored at -20°C.

**Control Antigen Reconstitution:** 100 µl water.

**Control Antigen Storage after Reconstitution:** -20 °C.

**Preadsorption Control:** 1 µg peptide per 1 µg antibody.


**Standard Quality Control of Each Lot:** Western blotting.

### Applications:

**Western Blotting:** Rat brain membranes (1:200).

**For research purposes only, not for human use.**

**Last Update:** May 15, 2005.



Western blotting of rat brain membranes:

1. Anti- K<sub>Ca</sub>2.3 (C-term) antibody (1:200).
2. Anti- K<sub>Ca</sub>2.3 (C-term) antibody, preincubated with the control peptide antigen.